

Table S1 - Comparison of patients included and not included in the study at 360 days.

Variable		Eligible n=454	Included n=189	Not included n=265	p*
		Mean ± standard deviation			
Age (years)		57.7 (14.5)	59.3 (13.4)	56.5 (15.3)	0.051
BMI§ (kg/m²)		31.0 (6.8)	31.3 (6.7)	30.6 (6.9)	0.370
Variable	Category				p*
		n (%)			
Center of study‡	Hospital das Clínicas	74 (16.3)	36 (19.0)	38 (14.4)	0.369
	Hospital Eduardo de Menezes	209 (46.1)	82 (43.4)	127 (48.1)	
	Hospital Júlia Kubistchek	170 (37.5)	71 (37.6)	99 (37.5)	
Sex‡	Female	233 (51.3)	96 (50.8)	157 (31.7)	0.849
	Male	221 (48.7)	93 (49.2)	128 (48.3)	
Hospitalization in ICU‡	No	200 (44.3)	95 (50.3)	105 (40.1)	0.032
	Yes	251 (55.7)	94 (49.7)	157 (59.9)	

‡ Missing data: ≤ 10%; § missing data: 56% (included n=147, not included n=107, eligible n=254). * ANOVA test to test differences between means and Pearson's Chi-square to test differences between proportions. BMI: body mass index. ICU: intensive care unit.